



Keith Corpus, MD

**Total Shoulder Arthroplasty**

This information and protocol is a guideline, individual variations and changes will apply

<u>Postop</u>	<u>Goals</u>	<u>Precautions</u>	<u>Exercises</u>
<b>Weeks 0-6</b> PT 1-2x/week HEP daily	Edema and pain control Protect subscap repair  Week 1: FF 100°, ER 0° Week 2: FF 120°, ER 10° Week 6: FF 150°, ER 20°	Sling when not doing exercises Limit ER to neutral x2 weeks No ER >20° (weeks 2-6) No active IR No backward extension (BE) No scapular retractions Limit abduction 75° No resisted elbow flexion	Elbow, wrist, hand ROM Codman/Pendulum Passive supine forward flexion Scapular mobility and stability (side-lying) Deltoid isometrics Posture training
<b>Weeks 6-12</b> PT 2-3x/week HEP daily	Protect subscap repair FF 160°, ER 45°	DC sling No resisted IR/BE No resisted scapular retractions Avoid painful ADL's	Advance P/AA/AROM Cane/pulley Rhythmic stabilization at 120° Begin AA→AROM IR/BE Submaximal isometrics ER/FF/ABD Closed chain kinetic exercises Scapular stabilization
<b>Weeks 12-16</b> PT 1-2x/week HEP daily	FF 160°, ER 60°, IR T12 Scapulohumeral rhythm UE strength 4/5	No sling Avoid painful ADL's Avoid activities that encourage scapula hiking or poor mechanics Limit strengthening to 3x/week to avoid rotator cuff tendinitis	Progress ROM/flexibility exercises Advance strengthening as tolerated Rhythmic humeral head stabilization Begin resisted IR/BE (isometrics→light bands→weights) Increase end ROM with passive stretch Begin eccentrics, plyometrics, and closed chain exercises when appropriate
<b>Weeks 16+</b> HEP daily	Maximal ROM Independent HEP	None	Progress strengthening, flexibility, and endurance